arXiv:1402.3836v2 [nlin.SI] 2 Mar 2014

Integrable Dispersive Chains and Energy Dependent Schrödinger Operator

Maxim V. Pavlov[1,2]
[1]Sector of Mathematical Physics,
Lebedev Physical Institute of Russian Academy of Sciences,
Leninskij Prospekt 53, 119991 Moscow, Russia
[2]Laboratory of Geometric Methods in Mathematical Physics,
Lomonosov Moscow State University,
Leninskie Gory 1, 119991 Moscow, Russia

Abstract

In this paper we consider integrable dispersive chains associated with the so called "Energy Dependent" Schrödinger operator. In a general case multi component reductions of these dispersive chains are new integrable systems, which are characterised by two arbitrary natural numbers. Also we show that integrable three dimensional linearly degenerate quasilinear equations of a second order possess infinitely many differential constraints. Corresponding dispersive reductions are integrable systems associated with the "Energy Dependent" Schrödinger operator.
MSC: 35F50, 35L65, 35Q53, 37K05, 37K10.

Keywords: commuting flow, conservation law, integrable system, Hamiltonian structure, reduction, quasilinear equation, dispersionless limit.

on the occasion of the 77th birthday of Professor Alexey Borisovich Shabat

Contents

1 Introduction **2**

2 Dispersive Integrable Chains **4**
2.1 Higher Commuting flows 4
2.2 Conservation Laws 6
2.3 Local Hamiltonian Structures 7

3 Multi Component Reductions **8**
3.1 Elementary Reductions 8
3.2 Rational Constraints with Movable Singularities 10
3.3 Negative Flows 11

4 The Exceptional Case **12**

5 Three Dimensional Linearly Degenerate Quasilinear Equations **13**
5.1 Negative Flows 16

6 The Dispersionless Limit **19**

7 Conclusion **22**

References **23**

1 Introduction

The remarkable Korteweg de Vries equation is associated with the linear Schrödinger equation

$$\psi_{xx} = (\lambda + u)\psi.$$

This paper is devoted to description of integrable systems associated with more general linear equation $\psi_{xx} = u\psi$, where a dependence $u(x, t, \lambda)$ with respect to the spectral parameter λ can be much more complicated (for instance, rational). According to [3] we call such a linear equation $\psi_{xx} = u(x, t, \lambda)\psi$ the energy dependent Schrödinger equation.

The function $\psi(x, t, \lambda)$ satisfies the pair of *linear* equations in partial derivatives

$$\psi_{xx} = u\psi, \quad \psi_t = a\psi_x - \frac{1}{2}a_x\psi. \tag{1}$$

Then the compatibility condition $(\psi_{xx})_t = (\psi_t)_{xx}$ yields the relationship

$$u_t = \left(-\frac{1}{2}\partial_x^3 + 2u\partial_x + u_x\right) a \tag{2}$$

between functions $u(x, t, \lambda)$ and $a(x, t, \lambda)$.

If we choose the linear dependences $u(x, t, \lambda) = \lambda + u^1(x, t)$ and $a(x, t, \lambda) = \lambda + a_1(x, t)$, we obtain nothing but the famous Korteweg de Vries equation[1]

$$u^1_t = \frac{1}{4}u^1_{xxx} - \frac{3}{2}u^1u^1_x, \tag{3}$$

where $a_1 = -\frac{1}{2}u^1$.

If we choose the quadratic dependence[2] $u(x, t, \lambda) = \lambda^2 + \lambda u^1(x, t) + u^2(x, t)$ and again the linear dependence $a(x, t, \lambda) = \lambda + a_1(x, t)$, we obtain nothing but the well-known Kaup–Boussinesq system

$$u^1_t = -u^1u^1_x + \left(-\frac{1}{2}u^1 + \xi\right)u^1_x + u^2_x, \quad u^2_t = \frac{1}{4}u^1_{xxx} - u^2u^1_x + \left(-\frac{1}{2}u^1 + \xi\right)u^2_x,$$

[1] An accurate derivation leads to $a_1 = -\frac{1}{2}u_1 + \xi$, where ξ is an arbitrary constant. However, corresponding equation $u_{1,t} = \frac{1}{4}u_{1,xxx} - \frac{3}{2}u_1u_{1,x} + \xi u_{1,x}$ reduces to the form (3) under the transformation $t \to t, x + \xi t \to x$.

[2] Here we follow S.J. Alber, see [2]

where $a_1 = -\frac{1}{2}u^1 + \xi$. The arbitrary constant ξ can be removed by a linear change of independent variables ($t \to t, x + \xi t \to x$) exactly as in the previous case. Thus finally the Kaup–Boussinesq system takes the form

$$u^1_t = u^2_x - \frac{3}{2}u^1 u^1_x, \quad u^2_t = \frac{1}{4}u^1_{xxx} - u^2 u^1_x - \frac{1}{2}u^1 u^2_x. \tag{4}$$

Multi-component *polynomial* (with respect to the spectral parameter λ) generalization $u(x,t,\lambda) = \lambda^M + \lambda^{M-1}u^1(x,t) + ... + u^M(x,t)$ was investigated in several papers [3]. Multi-component *rational* (with respect to the spectral parameter λ) generalization (ϵ_k are arbitrary parameters)

$$u(x,t,\lambda) = \frac{\lambda^M u^0(x,t) + \lambda^{M-1}u^1(x,t) + ... + u^M(x,t)}{\epsilon_M \lambda^M + \epsilon_{M-1}\lambda^{M-1} + ... + \epsilon_0} \tag{5}$$

was studied in [4]. The authors considered two main subclasses selected by the conditions: $\epsilon_M = 0$ and $u^0 = 1$ (the so called "Generalized KdV type systems"); $\epsilon_M = 0$ but $u^M = 1$ (the so called "Generalized Harry Dym type systems"). In another paper [14] we found a third narrow subclass determined by a sole restriction $u^M = 0$.

This paper is devoted to an open question: description of multi-component integrable systems associated with different dependencies $u(x,t,\lambda)$ with respect to the spectral parameter λ. In this paper we construct infinitely many multi component dispersive reductions of "integrable dispersive chains" introduced in [18]. Thus we found infinitely many new multi-component integrable systems. They can be written in a compact form (28), (29) and (41). Existence of dispersive reductions for linearly degenerate integrable hydrodynamic chains was discovered in [12]. The hierarchy of these hydrodynamic chains is associated with five integrable three dimensional quasilinear equations of a second order (see, for instance, [8]). First such a three dimensional quasilinear equation of a second order can be written in a hydrodynamic form (44). This equation possesses N component hydrodynamic reductions parameterised by N arbitrary functions of a single variable (see [15]). In this paper we show that integrable three dimensional quasilinear system of a first order (44) also possesses N component dispersive reductions, which are integrable dispersive systems (28), (47) characterised by two arbitrary natural numbers.

The structure of the paper is as follows. In Section 2 we considered general properties of integrable dispersive chains (associated with the energy dependent Schrödinger equation) like higher commuting flows, conservation laws and local Hamiltonian structures. In Section 3 we constructed new dispersive integrable systems. In Section 4 separately we presented an exceptional case with a dispersive chain but written in non-evolution form. In Section 5) we investigated three dimensional linearly degenerate quasilinear equations of a second order, we found explicit differential constraints which allows to reduce these quasilinear equations to dispersive integrable systems discussed in previous Sections. In Section 6 we briefly derived a dispersionless limit of integrable dispersive chains including their multi component dispersive reductions. In Section 7 we formulated a programme for a further research.

2 Dispersive Integrable Chains

According to [18], instead of the linear dependence $u(x,\mathbf{t},\lambda)=\lambda+u^1(x,\mathbf{t})$ (see (3)) and quadratic dependence $u(x,\mathbf{t},\lambda)=\lambda^2+\lambda u^1(x,\mathbf{t})+u^2(x,\mathbf{t})$ (see (4)) below we consider $(M=1,2,...)$

$$u(x,\mathbf{t},\lambda)=\lambda^M\left(1+\frac{u^1(x,\mathbf{t})}{\lambda}+\frac{u^2(x,\mathbf{t})}{\lambda^2}+\frac{u^3(x,\mathbf{t})}{\lambda^3}+...\right), \tag{6}$$

where u^k are infinitely many unknown functions[3].

The substitution (6) and the linear dependence $a^{(1)}=\lambda+a_1(x,\mathbf{t})$ into (2) yields Mth integrable dispersive chain[4]

$$u^k_t=u^{k+1}_x-\frac{1}{2}u^1u^k_x-u^ku^1_x+\frac{1}{4}\delta^k_Mu^1_{xxx},\quad k=1,2,..., \tag{7}$$

where δ^k_M is the Kronecker delta and

$$a_1=-\frac{1}{2}u^1. \tag{8}$$

2.1 Higher Commuting flows

Higher commuting flows of the Korteweg de Vries hierarchy are determined by the linear spectral system

$$\psi_{xx}=(\lambda+u^1)\psi,\quad \psi_{t^k}=a^{(k)}\psi_x-\frac{1}{2}a^{(k)}_x\psi,$$

where

$$a^{(k)}=\lambda^k+\sum_{m=1}^{k}a_m\lambda^{k-m}, \tag{9}$$

and functions a_m and u^1 depend on the "space" variable x and infinitely many extra "time" variables t^k (obviously, $t\equiv t^1$). Substitution (6) and (9) into (2) leads to higher commuting flows (here we also define $a_0=1$)

$$u^k_{t^s}=\sum_{m=0}^{s}\left(u^{k+m}\partial_x+\partial_xu^{k+m}-\frac{1}{2}\delta^{k+m}_M\partial^3_x\right)a_{s-m},\quad s=1,2,..., \tag{10}$$

where all coefficients a_m can be found iteratively from the linear system[5] (here we also define $u^0=1$ and $u^{-m}=0$ for all $m=1,2,...$)

$$\sum_{m=0}^{s}\left(u^{m-k}\partial_x+\partial_xu^{m-k}-\frac{1}{2}\delta^{m-k}_M\partial^3_x\right)a_{s-m}=0,\quad k=0,1,...,s-1. \tag{11}$$

[3] this Laurent expansion was suggested in [18]; see other detail in [1].

[4] earlier in [18] these integrable dispersive chains were written just in a compact symbolic form.

[5] Linear system (11) can be obtained formally from (10), if to fix $u^0=1$ and all other unknown functions $u^{-m}=0$ for $m=1,2,...$

For instance,

$$a_1 = -\frac{1}{2}u^1, \quad a_2 = -\frac{1}{2}u^2 + \frac{3}{8}(u^1)^2 - \frac{1}{8}\delta^1_M u^1_{xx},$$

$$a_3 = -\frac{1}{2}u^3 + \frac{3}{4}u^1u^2 - \frac{5}{16}(u^1)^3 + \frac{1}{32}\delta^1_M(10u^1u^1_{xx} + 5(u^1_x)^2 - u^1_{xxxx} - 4u^2_{xx}) - \frac{1}{8}\delta^2_M u^1_{xx}, ...$$

Thus all higher commuting flows are written in an evolution form. For instance, a first commuting flow to (7) is (here we identify $y \equiv t^2$)

$$u^k_y = u^{k+2}_x - \frac{1}{2}u^1u^{k+1}_x + \left(-\frac{1}{2}u^2 + \frac{3}{8}(u^1)^2 - \frac{1}{8}\delta^1_M u^1_{xx}\right)u^k_x - u^{k+1}u^1_x \tag{12}$$

$$+u^k\left(-u^2_x + \frac{3}{2}u^1u^1_x - \frac{1}{4}\delta^1_M u^1_{xxx}\right) + \frac{1}{4}\delta^{k+1}_M u^1_{xxx} + \frac{1}{4}\delta^k_M\left(u^2_{xxx} - \frac{3}{4}[(u^1)^2]_{xxx} + \frac{1}{4}\delta^1_M u^1_{xxxxx}\right).$$

A generating function of higher commuting flows is determined by the choice (instead of (9))

$$a = \frac{\mathbf{a}(x, \mathbf{t}, \zeta)}{\lambda - \zeta}, \tag{13}$$

where ζ is an arbitrary parameter. Indeed, substitution the asymptotic expansion ($\zeta \to \infty$)

$$\mathbf{a}(x, \mathbf{t}, \zeta) = -\zeta\left(1 + \frac{a_1(x, \mathbf{t})}{\zeta} + \frac{a_2(x, \mathbf{t})}{\zeta^2} + \frac{a_3(x, \mathbf{t})}{\zeta^3} + ...\right) \tag{14}$$

into (13) leads to

$$a = 1 + \frac{\lambda + a_1}{\zeta} + \frac{\lambda^2 + a_1\lambda + a_2}{\zeta^2} + \frac{\lambda^3 + a_1\lambda^2 + a_2\lambda + a_3}{\zeta^3} + ...,$$

where all polynomial expressions with respect to the spectral parameter λ are precisely (9). The substitution (6) and (13) into (2) implies (let us remind that $u^0 = 1$)

$$u^{k+1}_{\tau(\zeta)} = \zeta u^k_{\tau(\zeta)} + 2u^k\mathbf{a}_x + \mathbf{a}u^k_x - \frac{1}{2}\delta^k_M\mathbf{a}_{xxx}, \ k = 0, 1, ...,$$

where we introduced the special "time" variable $\tau(\zeta)$ instead of t in (2) to emphasize that we deal with a generating function of commuting flows. Taking into account that $u^1_{\tau(\zeta)} = 2\mathbf{a}_x$ and iteratively expressing all others $u^k_{\tau(\zeta)}$ via higher derivatives of u^m and $\mathbf{a}$, we come to a more explicit form

$$u^{k+1}_{\tau(\zeta)} = \sum_{m=0}^{k}\zeta^m(u^{k-m}\partial_x + \partial_x u^{k-m} - \frac{1}{2}\delta^{k-m}_M\partial^3_x)\mathbf{a}, \ k = 0, 1, ...,$$

which again yields (10), if to substitute (14) together with another formal expansion ($\zeta \to \infty$)

$$\partial_{\tau(\zeta)} = \partial_{t^0} + \frac{1}{\zeta}\partial_{t^1} + \frac{1}{\zeta^2}\partial_{t^2} + \frac{1}{\zeta^3}\partial_{t^3} + ...$$

Moreover, now we can identify $x \equiv t^0$.

2.2 Conservation Laws

The substitution

$$\psi = \exp\left(\int r dx\right)$$

into (1) yields

$$r_x + r^2 = u, \quad r_t = \left(ar - \frac{1}{2}a_x\right)_x. \tag{15}$$

Thus $r(x,t,\lambda)$ is a generating function of conservation law densities (with respect to the spectral parameter λ), while the second equation of (15) is a generating function of conservation laws.

The differential substitution (ϵ is an arbitrary constant)

$$r = \frac{\varphi_x}{2\varphi} + \frac{\epsilon}{2\varphi} \tag{16}$$

into the first equation of (15) yields

$$2\varphi\varphi_{xx} - \varphi_x^2 + \epsilon^2 = 4u\varphi^2, \tag{17}$$

which is nothing but a first integral of well-known equation (in the case of the Korteweg de Vries equation)

$$\varphi_{xxx} = 4u\varphi_x + 2u_x\varphi. \tag{18}$$

Indeed, the function $\varphi = \psi\psi^+$ satisfies two linear equations

$$\varphi_{xxx} = 4u\varphi_x + 2u_x\varphi, \quad \varphi_t = a\varphi_x - a_x\varphi, \tag{19}$$

where (see (1)) ψ and ψ^+ are two linearly independent solutions, which (obviously functionally dependent according to the Wronskian relationship $\psi\psi_x^+ - \psi^+\psi_x$ =const$\neq 0$) usually are determined by their asymptotic behavior ($\lambda \to \infty$): $\psi \to \exp(\lambda^{M/2}x), \psi^+ \to \exp(-\lambda^{M/2}x)$.

The second equation can be written in the conservative form

$$\left(\frac{1}{\varphi}\right)_t = \left(\frac{a}{\varphi}\right)_x. \tag{20}$$

Thus the function $p = 1/\varphi$ is a generating function of conservation law densities, and the above equation is a generating function of conservation laws. Taking into account (16), one can see that the second equation in (15) is equivalent to the above generating function of conservation laws (up to a total derivative).

Theorem: *Dispersive chains* (7), (36) *also have an alternative generation function of conservation laws*

$$(u'\varphi)_t = \left([2u + (\lambda + a_1)u']\varphi - \frac{1}{2}\varphi_{xx}\right)_x, \tag{21}$$

where $u' \equiv u'(\lambda)$, *i.e.*

$$u = \lambda^M + \sum_{k=1}^{\infty}\lambda^{M-k}u^k, \quad u' = M\lambda^{M-1} + \sum_{k=1}^{\infty}(M-k)\lambda^{M-k-1}u^k.$$

Proof: We seek a generating function of conservation law densities in the form $b\varphi$, where b is a *linear* expression with respect field variables u^k, whose coefficients depend on the spectral parameter λ only. Differentiating this product $b\varphi$ with respect to the independent variable "t" and taking into account that (see the first equation in (19)) $u_x\varphi = (2u\varphi - \frac{1}{2}\varphi_{xx})_x$, we expect that the flux of this generating function will be proportional to the function φ and its second derivative φ_{xx}. A straightforward computation yields that $b = u'$, the flux is $[2u + (\lambda + a_1)u']\varphi - \frac{1}{2}\varphi_{xx}$ and[6]

$$u'_t = (\lambda + a_1)u'_x + 2u'a_{1,x} + u_x,$$

where the function a_1 is defined by (8). The Theorem is proved.

Taking into account that (6) we seek an asymptotic expansion of (18) in the form (obviously, any solution of linear equation is determined up to an arbitrary factor; thus without loss of generality and for simplicity we choose the normalization to unity at infinity with respect to the spectral parameter λ)

$$\varphi(x, \mathbf{t}, \lambda) = 1 + \frac{\varphi_1(x, \mathbf{t})}{\lambda} + \frac{\varphi_2(x, \mathbf{t})}{\lambda^2} + \frac{\varphi_3(x, \mathbf{t})}{\lambda^3} + ..., \quad \lambda \to \infty. \tag{22}$$

Then we obtain (here we define $\varphi_0 = 1$ and remind that $u^0 = 1$)

$$\frac{1}{2}\sum_{k=0}^{\infty}\frac{\varphi_{k+1,xxx}}{\lambda^{M+k}} = \sum_{k=0}^{\infty}\frac{1}{\lambda^k}\sum_{m=1}^{k+1}(2u^{k+1-m}\varphi_{m,x} + \varphi_{k+1-m}u^m_x).$$

Selecting factors of each degree of the spectral parameter λ, one can see that expressions for coefficients φ_k coincide with expressions for coefficients a_k in (11), i.e. (22) becomes

$$\varphi(x, \mathbf{t}, \lambda) = 1 + \frac{a_1(x, \mathbf{t})}{\lambda} + \frac{a_2(x, \mathbf{t})}{\lambda^2} + \frac{a_3(x, \mathbf{t})}{\lambda^3} + ... \tag{23}$$

Thus substituting this asymptotic expansion into generating function (21), infinitely many conservation laws can be presented explicitly.

2.3 Local Hamiltonian Structures

Construction of local multi-Hamiltonian structures for polynomial and rational cases (5) was presented in [11], [3], [4].

A hierarchy of integrable dispersive chains (7) possesses *infinitely* many local Hamiltonian structures, first two of them are ($s = 1, 2, ...$)

$$u^k_{t^s} = \sum_{m=1}^{s+1}\left(u^{k+m-1}\partial_x + \partial_x u^{k+m-1} - \frac{1}{2}\delta_M^{k+m-1}\partial_x^3\right)\frac{\delta \mathbf{H}_{s+1}}{\delta u^m}, \ k = 1, 2, ...;$$

$$u^1_{t^s} = -2\partial_x\frac{\delta \mathbf{H}_{s+2}}{\delta u^1}, \quad u^k_{t^s} = \sum_{m=2}^{s+2}\left(u^{k+m-2}\partial_x + \partial_x u^{k+m-2} - \frac{1}{2}\delta_M^{k+m-2}\partial_x^3\right)\frac{\delta \mathbf{H}_{s+2}}{\delta u^m}, \ k > 1.$$

[6]This equation can be obtained directly from (2) by differentiation with respect to the spectral parameter λ.

Remark: The first local Hamiltonian structure follows from (10), where we utilized the observation

$$a_m = \frac{\delta \mathbf{H}_{m+s}}{\delta u^s}, \ m = 0, 1, ...; \ s = 1, 2, ... \tag{24}$$

This is alternative approach for construction of local polynomial conservation laws (cf. (15), (20), (21)). In such a case all Hamiltonians can be found from above variation derivatives, for instance

$$\mathbf{H}_1 = \int u^1 dx, \quad \mathbf{H}_2 = \int \left(u^2 - \frac{1}{4}(u^1)^2 \right) dx,$$

$$\mathbf{H}_3 = \int \left(u^3 - \frac{1}{2}u^1 u^2 + \frac{1}{8}(u^1)^3 + \frac{1}{16}\delta^1_M (u^1_x)^2 \right) dx,$$

$$\mathbf{H}_4 = \int \Big(u^4 - \frac{1}{2}u^1 u^3 - \frac{1}{4}(u^2)^2 + \frac{3}{8}(u^1)^2 u^2 - \frac{5}{64}(u^1)^4$$
$$+ \frac{1}{32}\delta^1_M \left(-5u^1 (u^1_x)^2 - \frac{1}{2}(u^1_{xx})^2 + 4u^1_x u^2_x \right) + \frac{1}{16}\delta^2_M (u^1_x)^2 \Big) dx, ...$$

All other higher local Hamiltonian structures can be constructed utilizing the relationship (24), i.e.

$$\frac{\delta \mathbf{H}_s}{\delta u^m} = \frac{\delta \mathbf{H}_{s+k}}{\delta u^{m+k}}, \quad m, s, k = 1, 2, ...$$

For instance, the third local Hamiltonian structure is ($s = 1, 2, ...$)

$$u^1_{t^s} = -2\partial_x \frac{\delta \mathbf{H}_{s+3}}{\delta u^2}, \quad u^2_{t^s} = -2\partial_x \frac{\delta \mathbf{H}_{s+3}}{\delta u^1} - \left(u^1 \partial_x + \partial_x u^1 - \frac{1}{2}\delta^1_M \partial^3_x \right) \frac{\delta \mathbf{H}_{s+3}}{\delta u^2},$$

$$u^k_{t^s} = \sum_{m=3}^{s+3} \left(u^{k+m-3}\partial_x + \partial_x u^{k+m-3} - \frac{1}{2}\delta^{k+m-3}_M \partial^3_x \right) \frac{\delta \mathbf{H}_{s+3}}{\delta u^m}, \ k > 2.$$

3 Multi Component Reductions

Theory of multi-component semi-Hamiltonian hydrodynamic reductions of integrable hydrodynamic chains was build in [10] and applied in several papers [6], [15] (see also [13], [17]). Corresponding theory of multi-component integrable *dispersive* reductions of integrable *dispersive* chains does not exist at this moment. Nevertheless infinitely many multi-component integrable dispersive systems extracted from (7) are presented in this Section.

3.1 Elementary Reductions

Obviously for any natural number $N \geqslant M$ the reduction $u^{N+1} = 0$ of Mth dispersive chain (7) leads to N component integrable dispersive systems:

1. $N = M = 1$, this is the Korteweg de Vries equation (3);

2. $N=2, M=1$, this is the Ito system

$$u^1_t = u^2_x - \frac{3}{2}u^1u^1_x + \frac{1}{4}u^1_{xxx}, \quad u^2_t = -\frac{1}{2}u^1u^2_x - u^2u^1_x; \tag{25}$$

3. $N>2, M=1$,

$$u^1_t = u^2_x - \frac{3}{2}u^1u^1_x + \frac{1}{4}u^1_{xxx},$$

$$u^k_t = u^{k+1}_x - \frac{1}{2}u^1u^k_x - u^ku^1_x, \quad k=2,...,N-1,$$

$$u^N_t = -\frac{1}{2}u^1u^N_x - u^Nu^1_x;$$

4. $N=M>1$, (if $N=M=2$, this is the Kaup–Boussinesq equation, see (4))

$$u^k_t = u^{k+1}_x - \frac{1}{2}u^1u^k_x - u^ku^1_x, \quad k=1,2,...,N-1,$$

$$u^N_t = -\frac{1}{2}u^1u^N_x - u^Nu^1_x + \frac{1}{4}u^1_{xxx};$$

5. $N=M+1, M>1$,

$$u^k_t = u^{k+1}_x - \frac{1}{2}u^1u^k_x - u^ku^1_x, \ k=1,2,...,N-2,$$

$$u^{N-1}_t = u^N_x - \frac{1}{2}u^1u^{N-1}_x - u^{N-1}u^1_x + \frac{1}{4}u^1_{xxx},$$

$$u^N_t = -\frac{1}{2}u^1u^N_x - u^Nu^1_x.$$

6. $N>M+1, M>1$,

$$u^k_t = u^{k+1}_x - \frac{1}{2}u^1u^k_x - u^ku^1_x, \ k=1,...,M,$$

$$u^M_t = u^{M+1}_x - \frac{1}{2}u^1u^M_x - u^Mu^1_x + \frac{1}{4}u^1_{xxx},$$

$$u^k_t = u^{k+1}_x - \frac{1}{2}u^1u^k_x - u^ku^1_x, \quad k=M+1,...,N-1,$$

$$u^N_t = -\frac{1}{2}u^1u^N_x - u^Nu^1_x.$$

3.2 Rational Constraints with Movable Singularities

Now we consider more complicated N component reductions ($M = 1, 2, ...$)

$$u(x, \mathbf{t}, \lambda) = \frac{\lambda^{M+K} + \lambda^{M+K-1} v_{M+K-1}(x, \mathbf{t}) + ... + \lambda v_1(x, \mathbf{t}) + v_0(x, \mathbf{t})}{\lambda^K + \lambda^{K-1} w_{K-1}(x, \mathbf{t}) + ... + \lambda w_1(x, \mathbf{t}) + w_0(x, \mathbf{t})}, \quad K = 0, 1, ... \tag{26}$$

Suppose for simplicity that all roots of these two polynomials are pairwise distinct. Then the substitution

$$u(x, \mathbf{t}, \lambda) = \frac{\prod_{m=1}^{M+K} (\lambda - s^m(x, \mathbf{t}))}{\prod_{k=1}^{K} (\lambda - r^k(x, \mathbf{t}))} \tag{27}$$

into (2) together with the linear dependence $a^{(1)} = \lambda + a_1(x, \mathbf{t})$ yields new multi-component integrable dispersive systems

$$r^k_t = (r^k + a_1) r^k_x, \quad s^i_t = (s^i + a_1) s^i_x + \frac{1}{2} \frac{\prod_{k=1}^{K} (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} a_{1,xxx}, \tag{28}$$

where

$$a_1 = \frac{1}{2} \left(\sum_{m=1}^{M+K} s^m - \sum_{k=1}^{K} r^k \right). \tag{29}$$

Remark: In the particular case r^k =const, the ansatz (26) (ϵ_k are symmetric functions with respect to r^k according to the Viète theorem)

$$u(x, \mathbf{t}, \lambda) = \frac{\lambda^{M+K} + \lambda^{M+K-1} v_{M+K-1}(x, \mathbf{t}) + ... + \lambda v_1(x, \mathbf{t}) + v_0(x, \mathbf{t})}{\lambda^K + \epsilon_{K-1} \lambda^{K-1} + ... + \epsilon_1 \lambda + \epsilon_0}$$

was investigated in [4]. Then $2K + M$ component system (28) reduces to the form

$$s^i_t = (s^i + a_1) s^i_x + \frac{1}{2} \frac{\prod_{k=1}^{K} (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} a_{1,xxx}, \ i = 1, 2, ..., M + K,$$

where r^k =const. Thus all integrable systems considered in [4] also can be written in the above symmetric form. For instance, the Kaup–Boussinesq system (4) becomes

$$s^1_t = \frac{1}{2}(3s^1 + s^2) s^1_x + \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \ s^2_t = \frac{1}{2}(s^1 + 3s^2) s^2_x - \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}; \tag{30}$$

the Ito system (25) takes the form

$$s^1_t = \frac{1}{2}(3s^1 + s^2) s^1_x + \frac{s^1 (s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \ s^2_t = \frac{1}{2}(s^1 + 3s^2) s^2_x - \frac{s^2 (s^1 + s^2)_{xxx}}{4(s^1 - s^2)}.$$

3.3 Negative Flows

Substitution the expansion ($\zeta \to 0$)

$$\mathbf{a}(x, \mathbf{t}, \zeta) = -a_{-1}(x, \mathbf{t}) - \zeta a_{-2}(x, \mathbf{t}) - \zeta^2 a_{-3}(x, \mathbf{t}) + ...$$

into (13) yields (cf. (14))

$$a = -\frac{a_{-1}}{\lambda} - \left(\frac{a_{-2}}{\lambda} + \frac{a_{-1}}{\lambda^2}\right)\zeta - \left(\frac{a_{-3}}{\lambda} + \frac{a_{-2}}{\lambda^2} + \frac{a_{-1}}{\lambda^3}\right)\zeta^2 - ...$$

Substitution this expansion and the rational ansatz (27) into (2) implies infinitely many lower (negative) flows of the integrable hierarchies, whose first nontrivial (positive) members are (28). For instance, the choice

$$a = -\frac{a_{-1}}{\lambda} \tag{31}$$

determines non-evolution system

$$r^k_{t^{-1}} = -\frac{a_{-1}}{r^k} r^k_x, \quad s^i_{t^{-1}} = -\frac{a_{-1}}{s^i} s^i_x - \frac{1}{2s^i} \frac{\prod_{k=1}^{K}(s^i - r^k)}{\prod_{m \neq i}(s^i - s^m)} a_{-1,xxx}, \tag{32}$$

where

$$a_{1,t^{-1}} = a_{-1,x}. \tag{33}$$

Taking into account (29), one can derive an ordinary differential equation ($\tilde{\xi}$ is an integration constant) on the function a_{-1}, i.e. if any $r^k \neq 0$, then

$$\frac{(-1)^M}{2}\left(\frac{a_{-1,xx}}{a_{-1}} - \frac{1}{2}\frac{a^2_{-1,x}}{a^2_{-1}}\right) + \frac{\tilde{\xi}}{a^2_{-1}} = \frac{\prod_{i=1}^{M+K} s^i}{\prod_{k=1}^{K} r^k}. \tag{34}$$

If, for instance, $r^1 = 0$, then non-evolution system (32) becomes the *evolution* system ($k = 2, 3, ..., K; i = 1, 2, ..., M + K$), where ($\bar{\xi}$ is an integration constant)

$$a_{-1} = \bar{\xi}\left(\frac{\prod_{k=2}^{K} r^k}{\prod_{m=1}^{M+K} s^m}\right)^{1/2}. \tag{35}$$

4 The Exceptional Case

The concept of Mth dispersive integrable chain can be extended to the case $M = 0$ (see (6)). Indeed, the compatibility condition $(\psi_{xx})_t = (\psi_t)_{xx}$ of the linear system

$$\psi_{xx} = \left(1 + \frac{u^1}{\lambda} + \frac{u^2}{\lambda^2} + ...\right)\psi, \quad \psi_t = (\lambda + a_1)\psi_x - \frac{1}{2}a_{1,x}\psi$$

yields the *zeroth* dispersive integrable chain

$$u^k_t = u^{k+1}_x + a_1 u^k_x + 2u^k a_{1,x}, \quad k = 1, 2, ..., \tag{36}$$

where (ξ is an arbitrary constant)

$$\frac{1}{2}a_{1,xx} - 2a_1 = u^1 + \xi. \tag{37}$$

Since the function a_1 cannot be expressed via the function u_1 and a finite number of its derivatives, this zeroth dispersive chain is not an evolution system (in comparison with dispersive chains (7) with $M > 0$).

For this particular case we omit consideration of conservation laws, commuting flows and Hamiltonian structures, because such an investigation can be made exactly as in previous Sections. Here we just mention most important non-evolution reductions:

1. $N = 1$, the Camassa–Holm equation:

$$u^1_t = a_1 u^1_x + 2u^1 a_{1,x}, \quad \frac{1}{2}a_{1,xx} - 2a_1 = u^1 + \xi; \tag{38}$$

2. $N > 1$, the multi-component generalization of Camassa–Holm equation:

$$u^k_t = u^{k+1}_x + a_1 u^k_x + 2u^k a_{1,x}, \quad k = 1, 2, ..., N-1, \tag{39}$$

$$u^N_t = a_1 u^N_x + 2u^N a_{1,x}, \quad \frac{1}{2}a_{1,xx} - 2a_1 = u^1 + \xi.$$

3. Suppose for simplicity that all roots of two polynomials in (26) are pairwise distinct. Then the substitution

$$u(x, \mathbf{t}, \lambda) = \frac{\prod\limits_{m=1}^{K}(\lambda - s^m(x, \mathbf{t}))}{\prod\limits_{k=1}^{K}(\lambda - r^k(x, \mathbf{t}))} \tag{40}$$

into (2) yields multi-component non-evolution systems (cf. (28))

$$r^k_t = (r^k + a_1)r^k_x, \quad s^i_t = (s^i + a_1)s^i_x + \frac{1}{2}\frac{\prod\limits_{k=1}^{K}(s^i - r^k)}{\prod\limits_{m\neq i}(s^i - s^m)}a_{1,xxx},$$

where a_1 is determined by (ξ is an arbitrary constant, see (37))

$$2a_1 - \frac{1}{2}a_{1,xx} = \sum_{m=1}^{K} s^m - \sum_{k=1}^{K} r^k - \xi. \tag{41}$$

In this case lower commuting flows are determined by (31) and (40). Then (2) leads to (cf. (32))

$$r^k_{t^{-1}} = -\frac{a_{-1}}{r^k} r^k_x, \quad s^i_{t^{-1}} = -\frac{a_{-1}}{s^i} s^i_x - \frac{1}{2s^i} \frac{\prod_{k=1}^{K}(s^i - r^k)}{\prod_{m\neq i}(s^i - s^m)} a_{-1,xxx}, \tag{42}$$

where the function a_{-1} satisfies (33). Taking into account (41), one can derive an ordinary differential equation ($\tilde{\xi}$ is an integration constant) on the function a_{-1}, i.e. if any $r^k \neq 0$, then (cf. (34))

$$\frac{1}{2}(-1)^M \left(\frac{a_{-1,xx}}{a_{-1}} - \frac{1}{2}\frac{a^2_{-1,x}}{a^2_{-1}} \right) + \frac{\tilde{\xi}}{a^2_{-1}} = \frac{\prod_{m=1}^{K} s^m}{\prod_{k=1}^{K} r^k}.$$

If, for instance, $r^1 = 0$, then non-evolution system (42) becomes the *evolution* system

$$r^k_{t^{-1}} = -\frac{a_{-1}}{r^k} r^k_x, \quad s^i_{t^{-1}} = -\frac{a_{-1}}{s^i} s^i_x - \frac{1}{2} \frac{\prod_{k=2}^{K}(s^i - r^k)}{\prod_{m\neq i}(s^i - s^m)} a_{-1,xxx}$$

($k = 2, 3, ..., K; i = 1, 2, ..., K$), where ($\bar{\xi}$ is an integration constant, cf. (35))

$$a_{-1} = \bar{\xi} \left(\frac{\prod_{k=2}^{K} r^k}{\prod_{m=1}^{K} s^m} \right)^{1/2}.$$

5 Three Dimensional Linearly Degenerate Quasilinear Equations

Hierarchies of integrable dispersive chains (7), (10) have generating functions of conservation laws

$$p_{t^k} = (a^{(k)} p)_x,$$

where $p = 1/\varphi$ (see (20)). The compatibility conditions $(p_{t^k})_{t^m} = (p_{t^m})_{t^k}$ must be fulfilled, because corresponding dispersive integrable chains (and their multi component reductions) commute to each other. For instance, consistency of two first such equations (we remind that $x = t^0, t = t^1, y = t^2$)

$$p_t = [(\lambda + a_1)p]_x, \quad p_y = [(\lambda^2 + a_1\lambda + a_2)p]_x \tag{43}$$

leads to the three dimensional quasilinear system

$$a_{1,t} = a_{2,x}, \quad a_1 a_{2,x} + a_{1,y} = a_2 a_{1,x} + a_{2,t}, \tag{44}$$

whose integrability by the method of hydrodynamic reductions was investigated in [15]. This three dimensional quasilinear system also belongs to the class of such integrable systems which called *linearly degenerate*. This means that any of these systems admits at least one N component two dimensional hydrodynamic reduction (N must be arbitrary) which is *linearly degenerate* (see detail in [16], [5]). Moreover in such a case all hydrodynamic reductions can be completely described (see, for instance, [15]). They are parameterised by N arbitrary functions of a single variable.

In this Section we formulate the following

Statement: *Three dimensional quasilinear system* (44) *possesses infinitely many differential constraints* $a_1(\mathbf{u}, \mathbf{u}_x, ...), a_2(\mathbf{u}, \mathbf{u}_x, ...)$, *where field variables* u^k *are solutions of dispersive integrable systems determined by linear spectral problem* (1) *and described in Sections* **3** *and* **4**.

Already in two first papers [12] a concept of differential constraints was introduced for the hydrodynamic chain

$$a_{k,t} = a_{k+1,x} + a_1 a_{k,x} - a_k a_{1,x}, \ k = 1, 2, ..., \tag{45}$$

which can be obtained from the second equation in (1), where one should substitute the linear ansatz $a^{(1)} = \lambda + a_1$ instead of a and formal expansion (23) instead of φ. L. Martinez Alonso and A.B. Shabat also considered such remarkable examples like the Korteweg de Vries equation and the nonlinear Schrödinger equation.

The first commuting flow to (45)

$$a_{k,y} = a_{k+2,x} + a_1 a_{k+1,x} - a_{k+1} a_{1,x} + a_2 a_{k,x} - a_k a_{2,x}, \ k = 1, 2, ... \tag{46}$$

can be obtained from the second equation in (1), where one should substitute the quadratic ansatz $a^{(2)} = \lambda^2 + a_1\lambda + a_2$ instead of a and formal expansion (23) instead of φ. Taking first two equations $a_{1,t} = a_{2,x}, a_{2,t} = a_{3,x} + a_1 a_{2,x} - a_2 a_{1,x}$ from (45), the first equation $a_{1,y} = a_{3,x}$ from (46), and excluding $a_{3,x}$, one can obtain three dimensional quasilinear system (44).

Remark: *Hydrodynamic* chain (45) under differential invertible polynomial substitutions $a_k(\mathbf{u}, \mathbf{u}_x, ...)$ from (11) transforms into *dispersive* chains (7), where M is any natural number. This means that any Mth dispersive chain is *equivalent* to another Kth dispersive chain by appropriate invertible differential substitutions (also including the exceptional case $M = 0$). However, corresponding differential reductions are not equivalent to each other.

Substitution the quadratic dependence $a^{(2)} = \lambda^2 + \lambda a_1 + a_2$ into (2) together with ansatz (27) yields first commuting flow to (28)

$$r^k_y = (a_2 + a_1 r^k + (r^k)^2) r^k_x,$$

$$s^i_y = (a_2 + a_1 s^i + (s^i)^2) s^i_x + \frac{1}{2} \frac{\prod_{k=1}^{K} (s^i - r^k)}{\prod_{m \neq i} (s^i - s^m)} (s^i a_{1,xxx} + a_{2,xxx}), \tag{47}$$

where a_1 is determined by (29) and

$$a_2 = \frac{1}{4} \sum_{m=1}^{M+K} (s^m)^2 - \frac{1}{4} \sum_{k=1}^{K} (r^k)^2 + \frac{1}{2} a_1^2 + \frac{1}{4} \delta^1_M a_{1,xx}. \tag{48}$$

Thus in this Section we selected dispersive constraints (29), (48) of three dimensional quasilinear system (44), where $M + K$ functions $s^i(x, t, y)$ and K functions $r^k(x, t, y)$ are solutions of $2K + M$ component commuting dispersive integrable systems (28) and (47).

Now we present first four important examples of finite component differential reductions of three dimensional quasilinear system (44).

1. The Korteweg de Vries equation ($N = M = 1$). In this case commuting dispersive chains (7) and (12) reduce to (see (3))

$$u^1_t = \left(\frac{1}{4} u^1_{xx} - \frac{3}{4} (u^1)^2 \right)_x,$$

and its first commuting flow

$$u^1_y = \left(\frac{5}{8} (u^1)^3 - \frac{5}{16} (u^1_x)^2 - \frac{5}{8} u^1 u^1_{xx} + \frac{1}{16} u^1_{xxxx} \right)_x.$$

Substitution (see (11))

$$a_1 = -\frac{1}{2} u^1, \quad a_2 = \frac{3}{8} (u^1)^2 - \frac{1}{8} u^1_{xx}$$

into three dimensional quasilinear system (44) leads to an identity.

2. The Ito system ($N = 2, M = 1$). In this case commuting dispersive chains (7) and (12) reduce to (see (25))

$$u^1_t = u^2_x - \frac{3}{2} u^1 u^1_x + \frac{1}{4} u^1_{xxx}, \quad u^2_t = -\frac{1}{2} u^1 u^2_x - u^2 u^1_x,$$

and its first commuting flow

$$u^1_y = -\frac{3}{2} u^1 u^2_x + \left(-\frac{3}{2} u^2 + \frac{15}{8} (u^1)^2 - \frac{1}{8} u^1_{xx} \right) u^1_x$$

$$-\frac{1}{4} u^1 u^1_{xxx} + \frac{1}{4} u^2_{xxx} - \frac{3}{16} [(u^1)^2]_{xxx} + \frac{1}{16} u^1_{xxxxx},$$

$$u^2_y = \left(-\frac{3}{2}u^2 + \frac{3}{8}(u^1)^2 - \frac{1}{8}u^1_{xx}\right)u^2_x + \frac{3}{2}u^2u^1u^1_x - \frac{1}{4}u^2u^1_{xxx}.$$

Substitution (see (11))

$$a_1 = -\frac{1}{2}u^1, \quad a_2 = -\frac{1}{2}u^2 + \frac{3}{8}(u^1)^2 - \frac{1}{8}u^1_{xx},$$

into three dimensional quasilinear system (44) leads to an identity.

3. The Kaup–Boussinesq equation ($N = M = 2$). In this case commuting dispersive chains (7) and (12) reduce to (see (4))

$$u^1_t = u^2_x - \frac{3}{2}u^1u^1_x, \quad u^2_t = \frac{1}{4}u^1_{xxx} - u^2u^1_x - \frac{1}{2}u^1u^2_x,$$

and its first commuting flow

$$u^1_y = -\frac{3}{2}u^1u^2_x - \frac{3}{2}u^2u^1_x + \frac{15}{8}(u^1)^2u^1_x + \frac{1}{4}u^1_{xxx},$$

$$u^2_y = -\frac{3}{2}u^2u^2_x + \frac{3}{8}(u^1)^2u^2_x + \frac{3}{2}u^2u^1u^1_x + \frac{1}{4}u^2_{xxx} - \frac{3}{16}[(u^1)^2]_{xxx}.$$

Substitution (see (11))

$$a_1 = -\frac{1}{2}u^1, \quad a_2 = -\frac{1}{2}u^2 + \frac{3}{8}(u^1)^2$$

into three dimensional quasilinear system (44) leads to an identity.

4. The Camassa–Holm equation ($N = 1, M = 0$). Substitution expressions (ξ_1, ξ_2 are arbitrary parameters)

$$a_1 = -\frac{1}{2}\left(1 - \frac{1}{4}\partial^2_x\right)^{-1}(u^1 + \xi_1),$$

$$a_2 = \left(1 - \frac{1}{4}\partial^2_x\right)^{-1}\left(-\frac{1}{4}a_1a_{1,xx} - \frac{1}{8}a^2_{1,x} + \frac{3}{2}a^2_1 + \xi_1a_1 + \xi_2\right)$$

into three dimensional quasilinear system (44) leads to an identity, if the function $u^1(x, t, y)$ is a solution of the pair of nonlocal equations (see (38))

$$u^1_t = a_1u^1_x + 2u^1a_{1,x}, \quad u^1_y = 2u^1a_{2,x} + a_2u^1_x.$$

5.1 Negative Flows

Now we consider another asymptotic expansion of the function (cf. (13), (14))

$$\mathbf{a}(x, \mathbf{t}, \zeta) = -a_{-1} - \zeta a_{-2} - \zeta^2 a_{-3} - ..., \quad \zeta \to 0 \tag{49}$$

Then consistency of two generating functions of conservation laws (cf. (43))

$$p_{t^{-1}} = -\frac{1}{\lambda}(a_{-1}p)_x, \quad p_t = [(\lambda + a_1)p]_x$$

implies the second three dimensional quasilinear system (cf. (44))

$$a_{1,t^{-1}} = a_{-1,x}, \quad a_{-1,t} = a_1 a_{-1,x} - a_{-1} a_{1,x}; \tag{50}$$

consistency of two generating functions of conservation laws

$$p_{t^{-1}} = -\left(\frac{a_{-1}}{\lambda} p\right)_x, \quad p_{t^{-2}} = -\left[\left(\frac{a_{-2}}{\lambda} + \frac{a_{-1}}{\lambda^2}\right) p\right]_x \tag{51}$$

leads to the third three dimensional quasilinear system

$$a_{-1,t^{-2}} = a_{-2,t^{-1}}, \quad a_{-1,t^{-1}} = a_{-2} a_{-1,x} - a_{-1} a_{-2,x}. \tag{52}$$

Remark: Taking into account that $p = 1/\varphi$, substitution formal expansion (23) instead of φ into (51) yields two commuting hydrodynamic chains[7] to (45), (46)

$$a_{k+1,t^{-1}} = a_k a_{-1,x} - a_{-1} a_{k,x}, \ k = 0, 1, 2, ... \tag{53}$$

$$a_{k+2,t^{-2}} = a_{k+1} a_{-2,x} - a_{-2} a_{k+1,x} + a_k a_{-1,x} - a_{-1} a_{k,x}, \ k = -1, 0, 1, ... \tag{54}$$

Moreover, all above hydrodynamic chains can be extended on negative values of the index k (see, for instance, [15] and [13]). Thus, taking the first "negative" equation of hydrodynamic chain (45)

$$a_{-1,t} = a_1 a_{-1,x} - a_{-1} a_{1,x}$$

together with the first equation of hydrodynamic chain (53) $a_{1,t^{-1}} = a_{-1,x}$, one can obtain three dimensional quasilinear system (50); taking the first two "negative" equations of hydrodynamic chain (53)

$$a_{-1,t^{-1}} = a_{-2} a_{-1,x} - a_{-1} a_{-2,x}, \quad a_{-2,t^{-1}} = a_{-3} a_{-1,x} - a_{-1} a_{-3,x}$$

together with the first "negative" equation of hydrodynamic chain (54)

$$a_{-1,t^{-2}} = a_{-3} a_{-1,x} - a_{-1} a_{-3,x},$$

and eliminating the common block $a_{-3} a_{-1,x} - a_{-1} a_{-3,x}$, one can obtain three dimensional quasilinear system (52).

Instead of asymptotic expansion (49), we can consider an asymptotic expansion at any fixed point $\zeta = \lambda_0$. If we choose $\lambda_0 = 1$, then instead of (31), we obtain (see (13)) $a = b_{-1}(\lambda - 1)^{-1}$, where $b_{-1} = \mathbf{a}(x, \mathbf{t}, 1)$; if we keep λ_0 as an arbitrary constant, then we have another choice $a = c_{-1}(\lambda - \lambda_0)^{-1}$, where $c_{-1} = \mathbf{a}(x, \mathbf{t}, \lambda_0)$. Thus, we must introduce two extra independent variables y^{-1}, z^{-1} such that two extra generating functions of conservation laws

$$p_{y^{-1}} = -\left(\frac{b_{-1}}{\lambda - 1} p\right)_x, \quad p_{z^{-1}} = -\left(\frac{c_{-1}}{\lambda - \lambda_0} p\right)_x$$

can be determined as a part of an integrable hierarchy, which contains hydrodynamic chain (45) together with all its commuting flows. This means that all functions a_k (see,

[7] let us remind that $a_0 = 1$.

for instance, (45), (46), (49)), all functions u^k (see, for instance, (7)) and these three extra functions b_{-1}, c_{-1} and p depend on "time" variables t^k and y^{-1}, z^{-1} simultaneously. Consistency of two generating functions of conservation laws

$$p_{t^{-1}} = -\frac{1}{\lambda}(a_{-1}p)_x, \quad p_{y^{-1}} = -\left(\frac{b_{-1}}{\lambda - 1}p\right)_x$$

yields the fourth three dimensional quasilinear system

$$a_{-1,y^{-1}} = b_{-1,t^{-1}}, \quad a_{-1,y^{-1}} = b_{-1}a_{-1,x} - a_{-1}b_{-1,x}. \tag{55}$$

Remark: Introducing a potential function W such that $a_1 = W_x, a_2 = W_t, a_{-1} = W_{t^{-1}}, a_{-2} = W_{t^{-2}}, b_{-1} = W_{y^{-1}}$, systems (44), (50), (52), (55) can be written as four three dimensional quasilinear equations of a second order (let us remind that $x = t^0, t = t^1, y = t^2$), respectively

$$W_{t^0}W_{t^0t^1} + W_{t^0t^2} = W_{t^1}W_{t^0t^0} + W_{t^1t^1}, \quad W_{t^1t^{-1}} = W_{t^0}W_{t^0t^{-1}} - W_{t^{-1}}W_{t^0t^0}, \tag{56}$$

$$W_{t^{-1}t^{-1}} = W_{t^{-2}}W_{t^0t^{-1}} - W_{t^{-1}}W_{t^0t^{-2}}, \quad W_{t^{-1}y^{-1}} = W_{y^{-1}}W_{t^0t^{-1}} - W_{t^{-1}}W_{t^0y^{-1}}.$$

Now we consider three generating functions of conservation laws

$$p_{t^{-1}} = -\frac{1}{\lambda}(a_{-1}p)_x, \quad p_{y^{-1}} = -\frac{1}{\lambda - 1}(b_{-1}p)_x, \quad p_{z^{-1}} = -\frac{1}{\lambda - \lambda_0}(c_{-1}p)_x,$$

where $a_{-1} = W_{t^{-1}}, b_{-1} = W_{y^{-1}}$ and $c_{-1} = W_{z^{-1}}$. One can introduce a potential function S such that

$$dS = pdx - \frac{1}{\lambda}a_{-1}pdt^{-1} - \frac{b_{-1}}{\lambda - 1}pdy^{-1} - \frac{c_{-1}}{\lambda - \lambda_0}pdz^{-1}.$$

Then the compatibility conditions $(S_{t^{-1}})_{y^{-1}} = (S_{y^{-1}})_{t^{-1}}$ and $(S_{t^{-1}})_{z^{-1}} = (S_{z^{-1}})_{t^{-1}}$ yield

$$q_{y^{-1}} = \left(\frac{\lambda}{\lambda - 1}\frac{b_{-1}}{a_{-1}}q\right)_{t^{-1}}, \quad q_{z^{-1}} = \left(\frac{\lambda}{\lambda - \lambda_0}\frac{c_{-1}}{a_{-1}}q\right)_{t^{-1}},$$

where $q = a_{-1}p/\lambda$. The compatibility condition $(S_{z^{-1}})_{y^{-1}} = (S_{y^{-1}})_{z^{-1}}$ is equivalent to the compatibility condition $(q_{z^{-1}})_{y^{-1}} = (q_{y^{-1}})_{z^{-1}}$, which implies the fifth three dimensional quasilinear system

$$(\lambda_0 - 1)W_{t^{-1}}W_{y^{-1}z^{-1}} + W_{z^{-1}}W_{y^{-1}t^{-1}} - \lambda_0 W_{y^{-1}}W_{z^{-1}t^{-1}} = 0, \tag{57}$$

Recently a complete classification of linearly degenerate three dimensional quasilinear equations of a second order was presented in [8]. The list of these equations[8] coincides with five equations (56), (57).

Since all these five equations belong to the same hierarchy (the function W is common), the computation of differential reductions of last four three dimensional quasilinear equations of a second order reduces to the problem of computation of higher (positive) and lower (negative) commuting flows of integrable dispersive systems (28). For instance,

[8] all these three dimensional quasilinear equations of a second order can be found, for instance, in [15] and [13]. Some other references are in [8].

second three dimensional quasilinear system (50) possesses infinitely many differential constraints $a_1(\mathbf{u}, \mathbf{u}_x, ...), a_{-1}(\mathbf{u}, \mathbf{u}_x, ...)$ (see expressions (29), (34), (35)), where field variables $u^k(x, t, t^{-1})$ are solutions of dispersive systems (28) and (32). Moreover all lower differential constraints $a_{-k}(\mathbf{u}, \mathbf{u}_x, ...)$ can be found directly from (17) by substitution the expansion (cf. (23))

$$\varphi(x, \mathbf{t}, \lambda) = a_{-1} + \lambda a_{-2} + \lambda^2 a_{-3} + ..., \quad \lambda \to 0,$$

together with (27). Indeed, the ordinary differential equation on the function a_{-1} coincides with (34). The next coefficient a_{-2} is determined by the ordinary differential equation[9]

$$(-1)^M \frac{\prod\limits_{m=1}^{M+K} s^m}{\prod\limits_{k=1}^{K} r^k} \left(\sum_{p=1}^{K} \frac{1}{r^p} - \sum_{p=1}^{M+K} \frac{1}{s^p} \right) = \frac{a_{-2,xx}}{2a_{-1}} - \frac{a_{-1,x}}{2a_{-1}^2} a_{-2,x} \tag{58}$$

$$+ \left(\frac{a_{-1,x}^2}{2a_{-1}^3} - \frac{a_{-1,xx}}{2a_{-1}^2} - \frac{2\xi_0}{4a_{-1}^3} \right) a_{-2} + \frac{\xi_1}{4a_{-1}^2},$$

where the function a_{-1} as already determined by (34). Thus third three dimensional quasilinear system (52) possesses differential reductions (32) and

$$r^k_{t^{-2}} = -\left(\frac{a_{-2}}{r^k} + \frac{a_{-1}}{(r^k)^2} \right) r^k_x,$$

$$s^i_{t^{-2}} = -\left(\frac{a_{-2}}{s^i} + \frac{a_{-1}}{(s^i)^2} \right) s^i_x - \frac{1}{2(s^i)^2} \frac{\prod\limits_{k=1}^{K} (s^i - r^k)}{\prod\limits_{m \neq i} (s^i - s^m)} (s^i a_{-2,xxx} + a_{-1,xxx}),$$

where differential constraints $a_{-1}(\mathbf{u}, \mathbf{u}_x, ...), a_{-2}(\mathbf{u}, \mathbf{u}_x, ...)$ are determined by (34) and (58). More lengthy computations lead to similar but more complicated expressions for fourth three dimensional quasilinear system (55) and for the fifth three dimensional quasilinear equation of a second order (57). We omit corresponding derivations here.

6 The Dispersionless Limit

Hydrodynamic chain (45) also can be written in the form (see formula (53) in [15])

$$u^k_t = u^{k+1}_x - \frac{1}{2} u^1 u^k_x - u^k u^1_x. \tag{59}$$

[9] constants ξ_0, ξ_1 are first coefficients of the expansion $\epsilon^2(\lambda) = \xi_0 + \lambda \xi_1 + ...$ (see (17)). Obviously, the coefficient $\tilde{\xi}$ coincides with ξ_0.

Indeed, the substitution $\varphi = q^{-1/2}$ into the second equation in (19) yields

$$q_t = \left(\lambda - \frac{1}{2}u^1\right) q_x - q u^1_x,$$

where $a_1 = -u^1/2$ and (taking into account (23))

$$q = 1 + \frac{u^1}{\lambda} + \frac{u^2}{\lambda^2} + \frac{u^3}{\lambda^3} + ...$$

Thus hydrodynamic chain (59) is a dispersionless limit of Mth dispersive chain (7) for any $M > 0$ (also this statement is valid for $M = 0$, see Section 4). So, we come to an interesting observation: hydrodynamic chain (45) under infinitely many triangular invertible transformations (11) can be written as dispersive chain (7) (by infinitely many different ways depended on the natural number M), whose dispersionless limit (59) is equivalent to original hydrodynamic chain (45) up to invertible triangular point transformations. Hydrodynamic chain (59) and its higher commuting flows[10] have infinitely many local Hamiltonian structures. They can be easily obtain by dispersionless limit from local Hamiltonian structures of Mth dispersive chain (see Subsection 2.3). First three of them are ($s = 1, 2, ...$)

$$u^k_{t^s} = \sum_{m=1}^{s+1} (u^{k+m-1}\partial_x + \partial_x u^{k+m-1}) \frac{\partial h_{s+1}}{\partial u^m}, \ k = 1, 2, ...;$$

$$u^1_{t^s} = -2\partial_x \frac{\partial h_{s+2}}{\partial u^1}, \quad u^k_{t^s} = \sum_{m=2}^{s+2} (u^{k+m-2}\partial_x + \partial_x u^{k+m-2}) \frac{\partial h_{s+2}}{\partial u^m}, \ k = 2, 3, ...;$$

$$u^1_{t^s} = -2\partial_x \frac{\partial h_{s+3}}{\partial u^2}, \quad u^2_{t^s} = -2\partial_x \frac{\partial h_{s+3}}{\partial u^1} - (u^1\partial_x + \partial_x u^1) \frac{\partial h_{s+3}}{\partial u^2},$$

$$u^k_{t^s} = \sum_{m=3}^{s+3} (u^{k+m-3}\partial_x + \partial_x u^{k+m-3}) \frac{\partial h_{s+3}}{\partial u^m}, \ k = 3, 4, ...$$

All Hamiltonian densities $h_k(\mathbf{u})$ can be found from (43) by substitution $p = 1/\varphi$, where expansion of φ is determined by (23). For instance first conservation law densities are (corresponding conservation laws follow from (43) and (59), see other detail in [15])

$$h_1 = u^1, \quad h_2 = u^2 - \frac{1}{4}(u^1)^2, \quad h_3 = u^3 - \frac{1}{2}u^1u^2 + \frac{1}{8}(u^1)^3,$$

$$h_4 = u^4 - \frac{1}{2}u^1u^3 - \frac{1}{4}(u^2)^2 + \frac{3}{8}(u^1)^2u^2 - \frac{5}{64}(u^1)^4, ...$$

A dispersionless limit of dispersive reductions associated with ansatz (5) was investigated in [9]. In this Section we consider a dispersionless limit of dispersive reductions (28), i.e. the dispersionless systems

$$r^k_t = (r^k + a_1) r^k_x, \quad s^i_t = (s^i + a_1) s^i_x,$$

[10] these commuting flows are dispersionless limit of (10). Here coefficients (see (24)) $a_m = \frac{\partial h_{m+s}}{\partial u^s}, m = 0, 1, ..., s = 1, 2, ...$, where Hamiltonian densities $h_k(\mathbf{u})$ are determined below.

where

$$a_1 = \frac{1}{2}\left(\sum_{m=1}^{M+K} s^m - \sum_{k=1}^{K} r^k\right).$$

Thus the field variables r^k, s^m are Riemann invariants (cf. formula (6) in [9]). These coordinates r^k, s^m are most convenient for comparison of distinct integrable systems. For instance, Kaup–Boussinesq system (4), Ito system (25) and two component generalization of the Camassa–Holm equation (39) have the same dispersionless limit (see the end of Subsection 3.2)

$$s^1_t = \frac{1}{2}(3s^1 + s^2)s^1_x, \qquad s^2_t = \frac{1}{2}(s^1 + 3s^2)s^2_x.$$

However, these three integrable dispersive systems do not connected by any differential substitutions. They are different, but have the same dispersionless limit.

In paper [15] we proved that hydrodynamic chain (45) possesses N component hydrodynamic reductions

$$r^i_t = (r^i + a_1)r^i_x, \tag{60}$$

where ($f_k(r^k)$ are arbitrary functions)

$$a_1 = \sum_{m=1}^{N} f_m(r^m).$$

Thus, in this paper we proved that hydrodynamic type system (60) has a dispersive integrable extension (28) if

$$a_1 = \frac{1}{2}\sum_{m=1}^{N} \epsilon_m r^m,$$

where $\epsilon_k = \pm 1$, while in [9] all constants $\epsilon_k = 1$.

Remark[11]: Kaup–Boussinesq system (4) can be written in the conservative form

$$u^1_t = \left(w - \frac{1}{2}(u^1)^2\right)_x, \quad w_t = \left(\frac{1}{4}u^1_{xx} - u^1 w\right)_x, \tag{61}$$

where

$$w = u^2 - \frac{1}{4}(u^1)^2.$$

Under the invertible substitution

$$w = \rho \pm \frac{1}{2}u^1_x,$$

(61) becomes the Kaup–Broer system

$$u^1_t = \left(\rho - \frac{1}{2}(u^1)^2 \pm \frac{1}{2}u^1_x\right)_x, \quad \rho_t = -\left(u^1\rho \pm \frac{1}{2}\rho_x\right)_x. \tag{62}$$

[11] this derivation of the relationship between the Kaup–Boussinesq system (4), the Kaup–Broer system (62) and the nonlinear Schrödinger equation (63) belongs to Solomon J. Alber.

Under the invertible substitution

$$v = u^1 \pm \frac{1}{2}\frac{\rho_x}{\rho},$$

(62) reduces to the Hasimoto form of the nonlinear Schrödinger equation

$$v_t = \left(\rho - \frac{v^2}{2} - \frac{\rho_{xx}}{4\rho} + \frac{\rho_x^2}{8\rho^2}\right)_x, \quad \rho_t = -(v\rho)_x. \tag{63}$$

Indeed, under the Madelung substitution

$$\psi = \sqrt{\rho}e^{\int v dx}, \quad \psi^+ = \sqrt{\rho}e^{-\int v dx},$$

(63) transforms into the nonlinear Schrödinger equation

$$\psi_t + \frac{1}{2}\psi_{xx} - \psi\psi^+\psi = 0, \quad \psi_t^+ - \frac{1}{2}\psi_{xx}^+ + \psi^+\psi\psi^+ = 0. \tag{64}$$

Thus nonlinear Schrödinger equation (64) is reducible to the form (30), where

$$\rho = -\frac{1}{4}(s^1 - s^2)^2 \pm \frac{1}{2}(s^1 + s^2)_x, \quad v = -s^1 - s^2 \pm \frac{1}{2}(\ln\rho)_x.$$

Also nonlinear Schrödinger equation (64) is well known in the complex form

$$i\psi_t - \frac{1}{2}\psi_{xx} - \psi\psi^+\psi = 0, \quad i\psi_t^+ + \frac{1}{2}\psi_{xx}^+ + \psi^+\psi\psi^+ = 0, \tag{65}$$

which follows from (64) by the complex change of independent variables $x \to ix, t \to it$. In such a case (65) can be written in the form (cf. (30))

$$s_t^1 = \frac{1}{2}(3s^1 + s^2)s_x^1 - \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \quad s_t^2 = \frac{1}{2}(s^1 + 3s^2)s_x^2 + \frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}.$$

Thus, all above systems possess the same dispersionless limit. The field variables s^1, s^2 are natural coordinates for the nonlinear Schrödinger equation. For investigation of shock waves ($\epsilon \to 0$) the nonlinear Schrödinger equation can be written in the form (by rescaling $\partial_x \to \epsilon\partial_x, \partial_t \to \epsilon\partial_t$)

$$s_t^1 = \frac{1}{2}(3s^1 + s^2)s_x^1 - \epsilon^2\frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}, \quad s_t^2 = \frac{1}{2}(s^1 + 3s^2)s_x^2 + \epsilon^2\frac{(s^1 + s^2)_{xxx}}{4(s^1 - s^2)}.$$

Such a consideration obviously is valid just if the difference $s^1 - s^2$ is not small.

7 Conclusion

In this paper we found new multi component integrable dispersive systems (28) associated with the energy dependent Schrödinger equation. We believe that the same approach based on construction of dispersive chains can be utilized for all others linear spectral

problems. Moreover we hope that each linear spectral problem can be associated with some three dimensional linearly degenerate quasilinear system of a first order. Thus we would like to establish a link between three dimensional linearly degenerate quasilinear systems of a first order and two dimensional dispersive integrable systems.

More general alternative approach for extraction of integrable dispersive systems from (2) is based on extension of method of hydrodynamic reductions (see detail in [7]) to integrable dispersive deformations. In this case functions a_1 and a_2 (see (44)) depend on N Riemann invariants r^k and can be written in the quasipolynomial form

$$a_1 = a_{10}(\mathbf{r}) + \epsilon b_{1k}(\mathbf{r}) r^k_x + \epsilon^2 (c_{1km}(\mathbf{r}) r^k_x r^m_x + c_{1k}(\mathbf{r}) r^k_{xx}) + ...,$$

$$a_2 = a_{20}(\mathbf{r}) + \epsilon b_{2k}(\mathbf{r}) r^k_x + \epsilon^2 (c_{2km}(\mathbf{r}) r^k_x r^m_x + c_{2k}(\mathbf{r}) r^k_{xx}) + ...,$$

where Riemann invariants $r^k(x,t,y)$ satisfy two commuting systems

$$r^i_t = \lambda^i(\mathbf{r}) r^i_x + \epsilon(\lambda^i_{km}(\mathbf{r}) r^k_x r^m_x + \lambda^i_k(\mathbf{r}) r^k_{xx}) + ... \tag{66}$$

$$r^i_y = \mu^i(\mathbf{r}) r^i_x + \epsilon(\mu^i_{km}(\mathbf{r}) r^k_x r^m_x + \mu^i_k(\mathbf{r}) r^k_{xx}) + ... \tag{67}$$

Functions $a_{10}(\mathbf{r}), b_{1k}(\mathbf{r}), \lambda^i(\mathbf{r}), ...$ are not yet determined. However, substitution of these expressions into (44) leads to (see detail in [15])

$$a_{10}(\mathbf{r}) = \sum_{m=1}^{N} \Psi_m(r^m), \quad a_{20}(\mathbf{r}) = \frac{1}{2} a^2_{10} + \sum_{m=1}^{N} \int f_m(r^m) \Psi_m(r^m) dr^m,$$

$$\lambda^i(\mathbf{r}) = f_i(r^i) + a_{10}, \quad \mu^i(\mathbf{r}) = f^2_i(r^i) + a_{10} f_i(r^i) + a_{20}.$$

All other coefficients $c_{1km}(\mathbf{r}), c_{1k}(\mathbf{r}), \lambda^i_{km}(\mathbf{r}), \lambda^i_k(\mathbf{r}), ...$ can be found iteratively for each degree of the parameter ϵ. Integrability of systems (66), (67) follows from (2). Corresponding equations

$$u_t = 2u a_{1,x} + (\lambda + a_1) u_x - \frac{1}{2} \epsilon^2 a_{1,xxx},$$

$$u_y = 2u(\lambda a_{1,x} + a_{2,x}) + (\lambda^2 + \lambda a_1 + a_2) u_x - \frac{1}{2} \epsilon^2 (\lambda a_{1,xxx} + a_{2,xxx})$$

determine a common function $u(x,t,y,\lambda)$, which one can look for in the form

$$u = u_0(\mathbf{r}, \lambda) + \epsilon u_{1k}(\mathbf{r}, \lambda) r^k_x + \epsilon^2 [u_{2k}(\mathbf{r}, \lambda) r^k_{xx} + u_{2km}(\mathbf{r}, \lambda) r^k_x r^m_x] + ...$$

This investigation should be made in a separate paper.

Acknowledgement

MVP's work was also partially supported by the RF Government grant #2010-220-01-077, ag. #11.G34.31.0005, by the grant of Presidium of RAS "Fundamental Problems of Nonlinear Dynamics" and by the RFBR grant 14-01-00389. MVP thanks E.V. Ferapontov and G.A. El for useful comments and important discussions.

References

[1] *V. E. Adler, A.B. Shabat,* Model equation of the theory of solitons, *Theor. Math. Phys.* **153** No. 1 (2007) 1373–1387.

[2] *S.J. Alber,* Hamiltonian systems on the Jacobi varieties. The geometry of Hamiltonian systems (Berkeley, CA, 1989) 23–32, Math. Sci. Res. Inst. Publ., 22, Springer, New York, 1991. *S.J. Alber,* Associated integrable systems, *J. Math. Phys.* **32** No. 4 (1991) 916-922.

[3] *M. Antonowicz, A. Fordy,* A family of completely integrable multi-Hamiltonian systems, *Phys. Letts. A* **122** (1987) 95–99. *M. Antonowicz, A. Fordy,* Coupled KdV equations with multi-Hamiltonian structures, *Physica* **28D** (1987) 345–57.

[4] *M. Antonowicz, A. Fordy,* Coupled Harry Dym equations with multi-Hamiltonian structures, *J. Phys. A* **21** (1988) L269–75. *M. Antonowicz, A. Fordy,* Factorisation of energy dependent Schrödinger operators: Miura maps and modified systems, *Comm. Math. Phys.*, **124** (1989) 465-486.

[5] *E.V. Ferapontov*, Integration of weakly-nonlinear hydrodynamic systems in Riemann invariants, *Phys. Lett. A* **158** (1991) 112-118.

[6] *E.V. Ferapontov, K.R. Khusnutdinova, D.G. Marshall, M.V. Pavlov*, Classification of Integrable Hydrodynamic chains associated with Kupershmidt's brackets, *J. Math. Phys.*, **47** (2006) 103507-103520. *E.V. Ferapontov, D.G. Marshall*, Differential-geometric approach to the integrability of hydrodynamic chains: the Haantjes tensor, *Mathematische Annalen*, **339** No. 1 (2007) 61-99.

[7] *E.V. Ferapontov, A. Moro*, Dispersive deformations of hydrodynamic reductions of (2 + 1)D dispersionless integrable systems, *J. Phys. A* **42** No. 3 (2009) 035211. *E.V. Ferapontov, A. Moro, V.S. Novikov*, Integrable equations in 2 + 1 dimensions: deformations of dispersionless limits, *J. Phys. A* **42** No. 34 (2009) 345205.

[8] *E.V. Ferapontov, J. Moss*, Linearly degenerate PDEs and quadratic line complexes, arXiv:1204.2777, submitted.

[9] *E.V. Ferapontov, M.V. Pavlov*, Quasiclassical limit of coupled KdV equations. Riemann invariants and multi-Hamiltonian structure, *Physica* **52D** No. 2–3 (1991) 211–219.

[10] *J. Gibbons, S.P. Tsarev*, Reductions of Benney's equations, *Phys. Lett. A*, **211** (1996) 19-24. *J. Gibbons, S.P. Tsarev*, Conformal maps and reductions of the Benney equations, *Phys. Lett. A*, **258** (1999) 263-270.

[11] *L. Martinez Alonso,* Schrödinger spectral problems with energy–dependent potentials as sources of nonlinear Hamiltonian evolution equations, *J. Math. Phys.* **21** (1980) 2342–2349.

[12] *L. Martinez Alonso, A.B. Shabat,* Energy-dependent potentials revisited: a universal hierarchy of hydrodynamic type, *Phys. Letts. A* **299** No. 4 (2002) 359–365. *L. Martinez Alonso, A.B. Shabat,* Towards a Theory of Differential Constraints of a Hydrodynamic Hierarchy, *J. Nonlin. Math. Phys.* **10** No. 2 (2003) 229–242.

[13] *L. Martinez Alonso, A.B. Shabat,* Hydrodynamic Reductions and Solutions of a Universal Hierarchy, *Theor. Math. Phys.* **140** No. 2 (2004) 1073–1085.

[14] *M. Marvan, M.V. Pavlov,* Energy dependent Schrödinger operator and a new subclass of integrable systems, submitted to SIGMA.

[15] *M.V. Pavlov,* Integrable hydrodynamic chains, *J. Math. Phys.* **44** No. 9 (2003) 4134-4156.

[16] *M.V. Pavlov,* Hamiltonian formalism of weakly nonlinear systems in hydrodynamics, *Theor. Math. Phys.* **73** (1987) 1242–1245.

[17] *M.V. Pavlov*, The Kupershmidt hydrodynamic chains and lattices, *IMRN* (2006) article ID 46987. *M.V. Pavlov*, Classification of integrable hydrodynamic chains and generating functions of conservation laws, *J. Phys. A: Math. Gen.*, (2006) 10803-10819. *M.V. Pavlov*, Algebro-geometric approach in the theory of integrable hydrodynamic type systems, *Comm. Math. Phys.*, **272** No. 2 (2007) 469-505. *M.V. Pavlov*, The Hamiltonian approach in the classification and the integrability of hydrodynamic chains, ArXiv: Nlin.SI/0603057. *M.V. Pavlov, S.A. Zykov*, Classification of integrable conservative hydrodynamic chains, submitted (2010) arXiv: 0912.4954.

[18] *A.B. Shabat,* Universal solitonic hierarchy, *J. Nonlin. Math. Phys.* **12** Supplement 1 (2005) 614–624. *A.B. Shabat,* Symmetric polynomials and conservation laws, *Vladikavkazskij matematicheskij zhurnal* **14** No. 4 (2012) 83–94 (in Russian).